

Recoil-induced lasing

J. Javaloyes¹, M. Perrin¹, G.L. Lippi¹, and A. Politi^{1,2}

¹ *Institut Non Linéaire de Nice, UMR 6618 CNRS,
Université de Nice-Sophia Antipolis,
1361 Route des Lucioles,
F-06560, Valbonne France*

² *Istituto Nazionale di Ottica Applicata,
L.go E. Fermi 6, 50125 Firenze, Italy*

(Dated: February 2, 2008)

The interaction of an atomic gas confined inside a cavity with a strong electromagnetic field is numerically and theoretically investigated in a regime where recoil effects are not negligible. The spontaneous appearance of a density grating (atomic bunching) accompanied by the onset of coherent backpropagation is found to be ruled by a continuous phase-transition. Numerical tests allow us to convincingly prove that the transition is steered by the appearance of a periodic atomic density modulation. Consideration of different experimental relaxation mechanisms induces us to analyze the problem in nearly analytic form, in the large detuning limit, using both a Vlasov approach and a Fokker-Planck description. The application of our predictions to recent experimental findings, reported in [Phys. Rev. Lett. **91**, 183601 (2003)], yields a semi-quantitative agreement with the observations.

PACS numbers: 42.50.Vk, 05.45.Xt, 05.65.+b., 42.65.Sf

I. INTRODUCTION

Recoil in the interaction between atoms and electromagnetic fields is almost exclusively associated with the idea that momentum transfer is a way of slowing down atoms to extremely low-temperatures [1]. However, non trivial consequences on light propagation have been uncovered as well. For instance, in the realm of single-atom effects, the so-called Recoil Induced Resonances (RIR) have been investigated both theoretically [2] and experimentally [3]. Moreover, it has been conjectured that atomic recoil can collectively give rise to coherent propagation through backreflection from a spontaneously generated density grating: the so called Collective Atomic Recoil Laser (CARL) [4]. Finally, connections between the two approaches have been investigated in [5].

The very first observation of spontaneous backward emission was made in a ring cavity experiment [6], where the very low “transverse temperature” might have been responsible for CARL amplification. More detailed experimental accounts have been reported in Refs. [7, 8], although it was not possible to establish whether the generation of the backward field was due to the spontaneous formation of a density grating, or viceversa. In fact, although it was later ascertained that recoil plays a prominent role [9] in the underlying physics [7, 8], an alternative interpretation based on the formation of a standard polarization grating could not be entirely ruled out [10]. Only recently has the first convincing evidence of CARL been given in a beautiful experiment performed on a sample of cold Rubidium atoms [11].

On the theoretical side, too, no final conclusion could be drawn on the CARL features, because the first model allowed only investigating the transient regime. Stationary states have been obtained for the first time by includ-

ing collisions with an external buffer gas [12]. By simulating a low-temperature (a few mK) sodium vapour in the detuning range previously considered in the (numerical) literature, a nonequilibrium phase transition leading to a stationary nonzero backward field (above a given threshold) was therein identified. However, such a collective phenomenon could not be linked to the onset of a density grating, but rather to the birth of a nontrivial polarization grating [12].

The overall scenario has been observed also in a more general model accounting for the input-field dynamics [13] and has been successively clarified in [14] with the help of the elimination of the atomic variables; indeed, this step rendered the derivation of an effective free-energy potential for the backward field amplitude possible, thereby clearly showing the existence of a phase transition.

In this paper we show numerically and theoretically that this latter model also accounts for the CARL and allows one to understand its onset in terms of a continuous phase transition characterized by two effective variables. The resulting estimate of the critical point is compatible with the experimental results reported in [11]. The results presented here provide the first substantial and systematic progress towards a proof of the existence, in stable form, of a collective interaction, which, predicted years ago [4], had so far proven to be elusive.

More precisely, we investigate the transition which gives rise, spontaneously – i.e., without an external probe field –, to a coherent electromagnetic field counterpropagating with respect to the injected one. We choose to consider the ring-cavity-based model introduced in [13, 15] for two reasons. On the one hand, the available single-pass model [4] has intrinsic shortcomings [16] which may be of little consequence in particular circumstances, but

which renders it nonetheless less attractive. On the other hand, the most recent experimental results [11], which can be considered at the present time as the first and only available observation of a true CARL system, have been obtained in a ring cavity. Hence, the choice of a bidirectional ring cavity model imposes itself.

A strong concern, which tainted all previous predictions and experimental attempts at studying this problem, regards the influence of external perturbations (collisions, finite transit time in the interaction volume, “optical friction”, etc.) on the collective dynamics. One can conjecture that their presence could destroy the phase transition or, at least, affect it so strongly as to shift it to an unphysical region of parameter values. Our purpose in this work is to investigate such mechanisms using different approaches and mathematical/numerical tools, to obtain a convincing picture. Section II introduces the model we used in the investigation and discusses, in subsection II A, the main physical relaxation mechanisms that may take place in an experiment. The appropriate modeling tools for analyzing the two main mechanisms are introduced here. Section III is devoted to a systematic numerical analysis which allows us to highlight the origin of the forces that give rise to a transition of the CARL-type – i.e., caused by the appearance of a spatial density grating – as opposed to the one recently predicted [12, 14], based on an unusual polarization grating and possibly responsible for high-temperature experimental observations [7, 8]. The numerical approach makes it possible to isolate the different source terms and identify in which range of temperature each of them is predominant. Analytical descriptions present the advantage of offering a better understanding of the physics driving the system. Since the full problem is obviously analytically unmanageable, it is useful to study some limiting situations. This is done in Section IV, where we consider the case of large detuning between fields and atomic resonance. In this spirit, we describe the perturbations introduced by transit broadening (subsection IV A) and show that a second-order transition appears which can be described by expanding the spatial inhomogeneities in Fourier modes. The first and most relevant mode coincides with the *bunching parameter*, previously introduced to quantify the density grating [4]. Losses introduced by an optical molasses are studied in subsection IV B with the help of a Fokker-Planck equation that can be analytically solved in the large dissipation limit. Predictions resulting from the preceding analyses are offered in Section V, together with a comparison with the recent experimental results [11]. Some conclusions and perspectives are presented in Section VI.

II. MODEL

The starting model involves four dimensionless variables (one being complex) accounting for the single atom dynamics plus two complex equations describing the

amplitude of forward (x_f) and backward (x_b) uniform fields [13]

$$\begin{aligned}\dot{\theta}_j &= p_j, \\ \dot{p}_j &= -\Re[(x_f e^{i\theta_j} - x_b e^{-i\theta_j})s_j^*] (|\Delta_a|/2G), \\ \dot{s}_j &= G(x_f e^{i\theta_j} + x_b e^{-i\theta_j})d_j - (\Gamma_\perp - i\Delta_a)s_j, \\ \dot{d}_j &= -G\Re[(x_f e^{i\theta_j} + x_b e^{-i\theta_j})s_j^*] - \Gamma_\parallel(d_j + 1), \\ \dot{x}_f &= -(1 + i\Delta_c)x_f + Y + \tilde{C}\langle se^{-i\theta} \rangle, \\ \dot{x}_b &= -(1 + i\Delta_c)x_b + \tilde{C}\langle se^{i\theta} \rangle,\end{aligned}\tag{1}$$

where $1 \leq j \leq N$ and the angular brackets denote an average over the atomic ensemble. Time and related parameters are expressed in units of the photon life-time inside the cavity, $1/\kappa$, namely: the decay rates of the atomic polarization ($\Gamma_\perp$) and population inversion ($\Gamma_\parallel$), the detuning Δ_a between input field and atomic frequencies, and the cavity detuning Δ_c (i.e. the distance from the nearest cavity resonance). Moreover, $G = \sqrt{m\kappa}|\Delta_a|/\hbar/k$ and $\tilde{C} = \alpha\Gamma_\perp L/(GT)$ are the two coupling constants (m is the atomic mass, k the field wavenumber, α the unsaturated absorption rate per unit length, L the length of the atomic medium, and T the cavity’s transmittivity). The amplitude Y of the injected field is scaled such that $Y^2 = 4\mathcal{D}^2\mathcal{P}/(AG^2T\kappa^2\hbar^2)$, where $\mathcal{D}$ is the atomic dipole moment, $\mathcal{P}$ is the input power, and A the beam surface, expressed in physical units. An equivalent scaling is used for the forward and backward intracavity components, x_f and x_b . Finally: (i) $\theta_j = kz_j$ is the scaled atomic position; (ii) the atomic momentum P_j is rescaled to $p_j = kP_j/(m\kappa)$; (iii) s_j and d_j are the atomic polarization and population inversion, the first being expressed in units of $\mathcal{D}$.

A. Relaxation mechanisms

While dissipation mechanisms affecting the internal degrees of freedom and the field amplitudes are accounted for by the above model, no losses acting on the atomic momentum are therein included. As a result, atoms are continuously accelerated by radiation pressure, and Eqs. (1) can thereby describe only the transient regime corresponding to the early stages of CARL amplification.

In actual experimental situations, relaxation processes may be induced by various mechanisms, the main ones being: (a) collisions with a buffer gas (those between optically active atoms can be neglected under all realistic circumstances), (b) finite residence time in the interaction volume (added to some relaxation occurring outside it), and (c) “viscous” losses (e.g., as in an optical molasses). Collisions with a buffer gas require fairly high pressures and have been often used in the investigation of nonlinear coupling between atoms and light (cf., e.g., [17, 18]). In a numerical scheme, this relaxation processes can be implemented as a coupling with a thermal bath, efficiently simulated with “molecular dy-

namics” [19] techniques. More precisely, a random sequence of inter-collision times is independently generated for each atom according to an exponential distribution with average value equal to $1/\gamma_c$. At each collision, the momentum of the colliding, j -th, atom is randomly reset according to the Gaussian distribution $Q_0(p_j) = \exp\{-p_j^2/(2T)\}/\sqrt{2\pi T}$ where $T = k_B \bar{T} k^2/(m\kappa^2)$ is the rescaled temperature of the buffer gas (k_B is Boltzmann’s constant, $\bar{T}$ is the temperature in physical units); moreover, the phase of the atomic polarization s_j is also reset to a value uniformly distributed in the whole range $[0, 2\pi]$, as $|s_j|$ remains unchanged. Notice that this algorithm amounts to implicitly assuming that active and buffer-gas atoms have equal masses.

The finite transit time (b) introduces effective losses and has certainly played the main role in several experiments [6, 7, 8]. In such a situation, the atoms spend a more or less small fraction of time inside the interaction volume, which they enter and exit at random positions and in a random state (both for the velocity and for the internal degrees of freedom). Outside the interaction region, the atoms suffer thermalization, either through collisions with other atoms (e.g., a low pressure buffer gas), with a container, or are simply renewed through “fresh” atoms coming from a beam; hence, the effective relaxation comes from the loss of “synchronized” atoms (both for the external and the internal degrees of freedom) compensated, in average, by the entrance into the interaction volume, of “random” atoms. This phenomenon can be reinterpreted as a “reset” in the external and internal degrees of freedom of the atom in question. At the microscopic level, one can still adopt a description based on the thermalization scheme discussed in (a), by modelling each random exit/entrance in the active volume as a “collision”; the main difference with the previous case is that all atomic variables have now to be reset. At the macroscopic level, the mathematical tool most suited for such a description is a Vlasov equation, where the probability distribution for the atomic position and momentum are dynamically investigated. These considerations are the basis for the discussion of Section IV A.

The third mechanism (c), recently employed in [11], is an optical molasses, added in the experiment to obtain steady state backward emission. Here, the “molasses field” slows down the atoms by the standard cooling action while introducing a “friction” which prevents the Doppler shift from growing so large as to render the interaction with the field entirely negligible. It has been shown [20] that under the assumption of a small saturation parameter of the molasses field, the momentum dynamics is well described by a Langevin equation (compare with the second of Eqs. (1)),

$$\dot{p}_j = \Re e[(x_b e^{-i\theta_j} - x_f e^{i\theta_j}) s_j^*] (|\Delta_a|/2G) - \gamma p_j + \eta_j, \quad (2)$$

where η_j is a standard white noise,

$$\begin{aligned} \langle \eta_j(t) \rangle &= 0, \\ \langle \eta_j(t_1) \eta_k(t_2) \rangle &= 2\gamma T \delta_{jk} \delta(t_1 - t_2), \end{aligned} \quad (3)$$

(δ_{jk} is the Kronecker symbol) accounting for the inevitable fluctuations that thermalize the atom to a finite temperature T . In Section IV B, we will study this model in the limit of large detuning, a condition well verified in the experiment [11].

III. NUMERICAL RESULTS

We now intend to explore the predictions of the model, Eqs. 1, from a purely numerical point of view to provide a starting point for the later analysis. In particular, we are interested in studying the existence of a *steady state bunching* in the presence of a backward propagating field. We remark that the observation of a stable value of bunching in a bidirectional ring cavity has never been reported. Previous observations, obtained without cavity [4], have shown a strongly oscillating, irregular behaviour for this variable. In the other kind of observed transition [12, 13], instead, the θ distribution remains flat (within the fluctuations imposed by the sample size) both below and above threshold. Hence, the mere numerical observation of a stable, steady bunching represents an important step towards generalizing the CARL predictions [4].

Such an observation is, however, not sufficient in itself. The question arises naturally, whether the backward field is originated by the growth of a nonhomogeneous longitudinal atomic distribution, or whether the spatial inhomogeneity results as a consequence of a preexisting optical standing wave, generated by other source terms. Although apparently obvious, this point is far from being trivial. Indeed, any standing wave is going to generate a whole hierarchy of gratings (atomic polarization and inversion, index of refraction and – if the atoms are free to move – atomic density), but one needs to identify the true source. In this system, a source term which gives rise to a collective state has already been identified in the non-trivial grating in the atomic polarization’s phase [12, 14], while a standard polarization modulation (without phase transition) has been proposed [10] as a possible mechanism for interpreting experimental observations [7, 8].

In order to characterize the appearance of a spatial inhomogeneity in the atomic distribution, we introduce the characteristic function of the position (z_j) distribution defined in terms of its Fourier modes

$$B_n := \frac{1}{N} \sum_{j=1}^N e^{2inkz_j}. \quad (4)$$

Introducing the normalised variable θ , the general definition Eq. (4) reads, with our normalisation,

$$B_n = \langle e^{2in\theta} \rangle. \quad (5)$$

The direct integration of model (1), using the same techniques as outlined in [12] (i.e. with thermalization induced by collisions), provides the results shown in Fig. 1.

Fig. 1a shows the appearance of several Fourier modes, which reach a steady state amplitude after the backward field has grown to its final value (Fig. 1b, solid line). We stress that the latter grows from zero spontaneously, without any *seed* being injected in the computation.

From Fig. 1a we remark that one Fourier component of the atomic spatial distribution significantly grows away from zero well before the backward field begins to deviate from its initial value. We will take this as a suggestion, to be later verified, that the bunching is responsible for the appearance of the counterpropagating field. The harmonic components, instead, seem to be generated by the interaction of the inhomogeneously distributed atoms and the optical standing wave generated by the two macroscopic counterpropagating fields, (x_f and x_b). In addition, we also see how the forward field (dashed line in Fig. 1b) is modified by the presence of the spatially modulated atomic distribution.

The physical picture described by this figure is the following: the cavity initially contains the atoms and no fields. At time $t = 0$, we start injecting the external component Y , collinear with the forward direction. During the transient preceding synchronisation, the atoms use part of the energy to adapt their positions and velocities, thus building up a non homogeneous spatial distribution. We remark that the total energy injected into the cavity (see dot-dashed line) is larger than that contained in the two field components at steady state ($t > 300$ time units). The fact that both fields, x_f and x_b , grow nearly to the same value implies that one cannot neglect the dynamics of the forward field. This dynamics maximizes the spatial modulation and *both* fields adapt themselves to the global coupling that ensues. It is also crucial to remark that, although the spatial atomic inhomogeneity appears to be responsible for the generation of the backward field, at time $t \approx 230$ time units, the B_1 component has reached only one third of its final value. Hence, the rest of the modulation (fundamental plus all the harmonics), results from the interaction with the optical standing wave. This point is significant, since it implies that the scattering from one field into the other is going to be quite symmetric: any increase in one component will imply a larger number of photons in one direction, hence a larger scattered contribution from that direction into the other. The residual difference between the forward (dashed line, Fig. 1a) and the backward field is to be attributed to the fact that the former receives an additional contribution from the external injection; hence the corresponding mode contains more photons at any time.

A more convincing illustration of the active role played by the density grating in the generation of the backward field, can be gained by decomposing the “force” field acting on the intensity $|x_b|^2$ of the backward field into two contributions

$$\frac{d|x_b|^2}{dt} = \mathcal{F}(x_f, x_b) = -2|x_b|^2 + \mathcal{F}_A(x_f, x_b), \quad (6)$$




FIG. 1: Fig. (a), shows the time evolution of amplitude of the first Fourier modes of the θ -distribution ($n = 1, 2, 3$ correspond to solid, dashed and dotted curves, respectively). Fig. (b) shows the rise of both amplitudes (solid and dashed line corresponds to $|x_b|$ and $|x_f|$). The dot-dashed line represents the injected field, Y . Parameter are: $\Delta_a = -30$, $\Gamma_\perp = 1$, $\Gamma_\parallel = 2$, $\tilde{C} = 7.1$, $\Delta_c = 0$, $G = 54$, $Y = 1.45$, $\gamma_c = 1/3$, $T = 5 \cdot 10^{-2}$.

where $\mathcal{F}_A(x_f, x_b) = 2\tilde{C}\text{Re}(\langle se^{i\theta} \rangle x_b^*)$. The first term on the r.h.s., due to cavity losses, has always a stabilizing effect, while $\mathcal{F}_A(x_f, x_b)$, which accounts for the atomic contribution, can be either stabilizing or destabilizing. In order to gain some insight on the resulting dynamics, we compare the force fields generated in the presence and in the absence of bunching, respectively. We do this under the assumption that the atomic degrees of freedom can be neglected [21], i.e. that the relevant dynamical properties are contained in the dependence of $\mathcal{F}_A$ on x_b and x_f . As long as the atomic variables rapidly converge to a stationary state, one can treat $|x_f|$ and $|x_b|$ as if they were fixed parameters and thereby numerically determine the “force” field for different choices of the two amplitudes. In principle, one should also take care of the time-dependence associated with the detuning ν between the two fields; however, numerical simulations of the full model have revealed that the “force” field depends very weakly on ν . The force resulting for the same parameter values as in Fig. 1 is plotted in Fig. 2 for different values of the backward field amplitude. The solid line with no symbols represents the total effective “force” acting on the backward field. Steady state operation exists only when the resulting total force is zero, and its stability is obviously determined by the local slope. It is clear that the zero-field state is unstable, while $|x_b| \approx 0.6$ is a stable one. In this latter regime, one can state that the destabilizing action of the atomic contribution, $\mathcal{F}_A$ (dashed curve without symbols in Fig. 2), is balanced by the stabilizing effect due to the cavity loss term, $|x_b|^2$.

In order to test of the *true role* played by atomic

bunching (i.e., as a *master* or a *slave* variable), we have proceeded to “switching off” both collisions and optical forces acting on the atoms (this is obviously possible only in a simulation). The corresponding curves representing $\mathcal{F}_A$ (dashed line in Fig. 2) and the total force, $\mathcal{F}$, (solid line) are identified by symbols. As a result of the absence of external perturbations, each atomic position θ_j evolves linearly according to the velocity value at the “switch off” time; in this way, the density grating is rapidly washed out. This procedure allows us to separate the possible gain component coming from the density grating from the one arising in a pumped ensemble of two-level systems [22]. The stationary value of $\mathcal{F}_A$ under such conditions remains always negative (so does *a fortiori* the total force), indicating that the presence of a density grating is an *indispensable* ingredient to maintain a finite backward field. Notice that this is at variance with the transition studied in [12], where the force field would be positive even in the absence of a density grating [14]. Hence, we can safely conclude that the onset of a finite backward field is a clear example of CARL and that the steady state bunching obtained is the *cause* for the appearance for the backward field component, and not a *consequence* of the existence of an optical standing wave (cf., e.g., [10]) in the parameter range which we are investigating.




FIG. 2: The force $\mathcal{F}_A$ (dashed lines) and $\mathcal{F}$ (solid lines) versus the backward field amplitude x_b . Curves with no symbols correspond to the simulation of the full model; diamonds display curves obtained by imposing a flat distribution of positions θ (same parameter values as in Fig. 1).

IV. LARGE DETUNING LIMIT

In Section III we have just reported purely numerical observations. In the perspective of constructing an analytical description of the transition, it is important to identify the possibly few relevant variables. For this purpose we concentrate on the large detuning limit, where the standard adiabatic elimination holds. Although this allows us to analytically treat only a limiting case, we will draw from it the benefit of later comparing the predictions of this approach to the recent experimental observations [11], obtained with very large detuning between

field and atoms ($\Delta_a \approx 10^6 \gg 1$). A discussion of the predictions resulting from such an analysis will be offered in Section V.

By setting, in Eqs. (1), $\dot{d}_j = \dot{s}_j = 0$, expanding the solutions in powers of $1/\Delta_a$ (to be considered as a smallness parameter) and considering first order terms, one finds $d_j = -1$ and

$$s_j = -\frac{iG}{\Delta_a}(x_f e^{i\theta_j} + x_b e^{-i\theta_j}) \quad . \quad (7)$$

Accordingly, our model, Eqs. (1), reduces to the following set of equations

$$\begin{aligned} \dot{\theta}_j &= p_j, \\ \dot{p}_j &= \Im m(x_f x_b^* e^{2i\theta_j}), \\ \dot{x}_f &= -[(1 + i\bar{\Delta})x_f - Y + iCx_b B_{-1}], \\ \dot{x}_b &= -[(1 + i\bar{\Delta})x_b + iCx_f B_1], \end{aligned} \quad (8)$$

where $C = \tilde{C}G/\Delta_a$, and $\bar{\Delta} = \Delta_c + C$ accounts for the combined effect of atomic and cavity frequency shifts in the absence of bunching (i.e. $B_1 = B_{-1} = 0$). The above form is similar to the one discussed in [23], which was, however, obtained by phenomenologically adding relaxation terms to a single pass model [24].

A. Vlasov Model

It is now convenient to pass to a Vlasov-like description by introducing the distribution $Q(\theta, p)$ of positions and momenta. In order to simplify the analysis we now assume that also the atomic positions are randomized in each collision [25]. Accordingly, we can write

$$\partial_t Q + p\partial_\theta Q + \Im m(x_f x_b^* e^{2i\theta})\partial_p Q = -\gamma_c(Q - Q_0), \quad (9)$$

where ∂_y denotes the partial derivative with respect to y . In the absence of a backward field, $Q(\theta, p)$ converges towards the equilibrium distribution Q_0 .

We now study the stability of the solution $x_f = Y/(1 + i\bar{\Delta}) := Y_0$, $x_b = 0$, and $Q = Q_0$, by introducing the perturbations δx_f , δx_b , δQ . Since the equation for x_f depends on the other variables only at second order in the perturbations, it can be solved separately. Moreover, δx_f affects δx_b only at second order, so that it can be factored out. Therefore, the stability of δx_f justifies the assumptions made in deriving the single-pass model where x_f is considered to be a constant parameter [4, 12, 14]. One is thus left with the following two equations

$$\dot{\delta x}_b = -(1 + i\bar{\Delta})\delta x_b - iC\bar{Y}\delta B_1, \quad (10)$$

$$\partial_t \delta Q = -\partial_p Q_0 \Im m(\bar{Y} \delta x_b^* e^{2i\theta}) - (\gamma_c + p\partial_\theta)\delta Q. \quad (11)$$

From the structure of such equations we see that if the bunching is washed out – i.e., $B_{\pm 1} = 0$ and $\delta Q = 0$ – no instability can be expected in this regime. This is at variance with the case discussed in [12, 14] where the

effective stability of x_b can be turned into an instability sufficiently close to the atomic resonance.

From Eq. (10), one realizes that the coupling occurs through the first Fourier mode of the density distribution. One can thus solve the above equation by introducing the following Ansätze,

$$\delta Q(\theta, p, t) = \mathcal{Q}(p)e^{-2i\theta + \lambda t} + c.c. \quad , \quad \delta x_b = \mathcal{E}_b e^{\lambda t}, \quad (12)$$

where λ is a complex stability eigenvalue. After substituting the above expressions into Eqs. (10, 11) and equating terms with the same exponential factor, one finds

$$(\lambda + 1 + i\bar{\Delta}) \mathcal{E}_b + 2i\pi C\bar{Y} \int_{-\infty}^{\infty} dp \mathcal{Q}(p) = 0, \quad (13)$$

$$\frac{i}{2} \bar{Y}^* \mathcal{E}_b \partial_p Q_0 + (\lambda - 2ip + \gamma_c) \mathcal{Q} = 0. \quad (14)$$

By solving Eq. (14) and replacing into Eq. (13), one obtains the solvability condition

$$\lambda = - (1 + i\bar{\Delta}) - \pi C |\bar{Y}|^2 \int_{-\infty}^{\infty} dp \frac{\partial_p Q_0}{\lambda - 2ip + \gamma_c}. \quad (15)$$

Within this framework, the onset of both bunching and backward field is signalled by a change of sign of the real part of λ , λ_r [26]. The integral in Eq. (15) can be analytically expressed using an error function. Nevertheless, this would hide an important point. Let place ourselves at threshold, and rewrite Eq. (15), assuming that λ is purely imaginary. Setting $\lambda = -i\beta$, we obtain for the real part

$$0 = -1 + \pi \frac{C\gamma_c |\bar{Y}|^2}{T\sqrt{2\pi T}} \int_{-\infty}^{\infty} dp \frac{p e^{-p^2/(2T)}}{(\beta + 2p)^2 + \gamma_c^2}. \quad (16)$$

In order for such an equation to be valid, the condition

$$\int_{-\infty}^{\infty} dp \frac{p e^{-p^2/(2T)}}{(\beta + 2p)^2 + \gamma_c^2} > 0 \quad (17)$$

must be fulfilled. The numerator is an odd function of p . The sign of the integral in Eq. (17) is thus determined by the denominator, whose minimum is reached at $p_o = -\beta/2$. Finally, Eq. (17) imposes $\beta < 0$. From the sign convention adopted here (the same that in Ref. [13]), a complex field E , of frequency ω , is linked to its slowly varying envelope A as $E = A \exp[-i\omega t]$. Thus β physically represents the detuning between the backward field and the forward field. One can thus conclude that the spontaneous symmetry breaking causing x_b to rise can only give rise to a red detuned field.

B. Fokker-Planck description

We now turn to analyzing the relaxation mechanism that most closely models the recent experimental results [11]. There, the cw superposition of additional

laser beams, tuned to the D₂ transition of Rubidium, introduces a mechanism which slows down those atoms that would be accelerated away from resonance by the strong interaction arising from CARL. Indeed, it was reported [11] that in the absence of the molasses only transient backwards emission could take place (and its interpretation as CARL is complicated by the technique used for preparing the sample, which pre-traps the atoms in a standing wave). The key point in the experiment is that the molasses is always present during the interaction but on the *other* D-line (the experiment being done on the D₁).

In the large detuning limit, the second of Eq. (8) is replaced by

$$\dot{p}_j = \Im m(x_f x_b^* e^{2i\theta_j}) - \gamma p_j + \eta_j(t). \quad (18)$$

Eq. (18) describes the evolution of (θ, p) in a potential self-consistently determined by the dynamics of the two counter-propagating field components, x_f and x_b , which in turn depends on the joint distribution $Q(\theta, p)$. The evolution of Q corresponding to Eq. (18), is given by the Fokker-Planck equation

$$\partial_t Q + p \partial_\theta Q + \Im m(x_f x_b^* e^{2i\theta}) \partial_p Q = \gamma \partial_p (pQ + T \partial_p Q). \quad (19)$$

Notice that this equation differs from the Vlasov equation (9) only for the relaxation terms contained in the right hand side.

By proceeding as in the previous subsection, we linearize Eq. (19) obtaining

$$\begin{aligned} \partial_t \delta Q = & - \partial_p Q_0 \Im m(\bar{Y} \delta x_b^* e^{2i\theta}) \\ & - p \partial_\theta \delta Q + \gamma \partial_p (p \delta Q + T \partial_p \delta Q), \end{aligned} \quad (20)$$

which is to be compared to the corresponding Eq. (11) within the Vlasov description (the linearised x_b equation, being identical to Eq. (10) is not reported). Using the Ansätze (12), we finally find

$$\begin{aligned} \frac{i}{2} \bar{Y}^* \mathcal{E}_b \partial_p Q_0 + (\lambda - 2ip - \gamma) \mathcal{Q} \\ - \gamma (p \partial_p \mathcal{Q} + T \partial_p^2 \mathcal{Q}) = 0, \end{aligned} \quad (21)$$

which is to be solved together with the first of Eqs. (21). In view of its differential structure, it is doubtful whether a general analytic solution can be found, and the development of a numerical analysis would be a quite delicate task as well. Nevertheless, in the “strong friction” (SF) limit, the atomic momentum can be adiabatically eliminated and one can thereby perform a quite detailed investigation of the transition scenario. While a careful discussion of the adiabatic elimination in the presence of noise can be found in Ref. [27], here we limit ourselves to presenting a sketchy but substantially correct derivation where we set $\dot{p} = 0$ in Eq. (18). By proceeding in this way, one obtains the Langevin equation

$$\dot{\theta}_j = \frac{1}{\gamma} \Im m(x_f x_b^* e^{2i\theta_j}) + \frac{\eta_j}{\gamma}, \quad (22)$$

that is equivalent to the one-variable Fokker-Planck (Smoluchowski) equation

$$\partial_t \rho + \frac{1}{\gamma} \partial_\theta \Im m (x_f x_b^* e^{2i\theta}) \rho - \frac{T}{\gamma} \partial_\theta^2 \rho = 0 \quad (23)$$

for the variable ρ

$$\rho(\theta) = \int_{-\infty}^{\infty} Q(\theta, p) dp. \quad (24)$$

We remark that Eq. (22), which describes the dynamics of an ensemble of mean field coupled oscillators, belongs to the class of Kuramoto systems [28]. The main differences are: a Dirac-like distribution of eigenfrequencies and a mean field self-consistently provided by a dynamical equation. By paralleling the approach in Section IV A, the stability of the solution $x_f = \bar{Y}$, $x_b = 0$, $\rho = 1/2\pi$ can be studied by introducing the infinitesimal perturbations δx_b and $\delta \rho$. The linearisation of Eq. (23) yields

$$\partial_t \delta \rho = -\frac{1}{2\pi\gamma} \partial_\theta \Im m (\bar{Y} \delta x_b^* e^{2i\theta}) + \frac{T}{\gamma} \partial_\theta^2 \delta \rho. \quad (25)$$

Upon then inserting the Ansätze $\delta \rho = r e^{-2i\theta + \lambda t} + c.c$ and $\delta x_b = \mathcal{E}_b e^{\lambda t}$ into the above equation, one obtains

$$r = -\frac{\mathcal{E}_b \bar{Y}^*}{2\pi} \frac{1}{\lambda\gamma + 4T}, \quad (26)$$

and from the linearised equation for the field dynamics, Eq. (10),

$$\lambda = -(1 + i\bar{\Delta}) + i \frac{C|\bar{Y}|^2}{\lambda\gamma + 4T}, \quad (27)$$

where now,

$$B_{\pm 1} = \int_0^{2\pi} e^{\pm 2i\theta} \rho(\theta) d\theta. \quad (28)$$

The threshold can be now obtained by setting $\lambda = -i\beta$ (where β is a real number) in Eqs. (26,27). The resulting two real equations for β and $|\bar{Y}|$ are

$$\gamma\beta^2 - \bar{\Delta}\gamma\beta - 4T = 0, \quad (29)$$

$$-\beta(\gamma + 4T) + 4\bar{\Delta}T = C|\bar{Y}|^2. \quad (30)$$

Only the negative β solution of Eq. (29) leads to a physically acceptable solution $|\bar{Y}|^2 > 0$. The threshold equations finally read

$$\beta_{TH} = \frac{\bar{\Delta}}{2} - \sqrt{\left(\frac{\bar{\Delta}}{2}\right)^2 + 4\frac{T}{\gamma}}, \quad (31)$$

$$|\bar{Y}_{TH}|^2 = \frac{\bar{\Delta}}{2C}(4T - \gamma) + \frac{1}{C}(4T + \gamma)\sqrt{\left(\frac{\bar{\Delta}}{2}\right)^2 + 4\frac{T}{\gamma}}.$$

The first equation shows that the probe field is *always* red detuned at threshold. This result is in agreement with

the outcome of the Vlasov model (see Section IV A) even though these two models correspond to different thermalisation mechanisms.

In the case of a resonant cavity field, $\bar{\Delta} = 0$, the above equations reduce to

$$\beta_{TH} = -2\sqrt{\frac{T}{\gamma}}, \quad (32)$$

$$|\bar{Y}_{TH}|^2 = \frac{2}{C}(4T + \gamma)\sqrt{\frac{T}{\gamma}}. \quad (33)$$

In the SF limit, one can not only determine analytically the threshold condition, but can also go beyond, describing the regime above threshold. By inserting the expansion of ρ ,

$$\rho(\theta, t) = \sum_{n=-\infty}^{\infty} f_n(t) e^{in(2\theta + \beta t)}, \quad (34)$$

into Eq. (23), one obtains the recurrence relation

$$\gamma \dot{f}_n = -in\gamma\beta f_n - n[Rf_{n-1} - R^*f_{n+1}] - 4Tn^2 f_n, \quad (35)$$

where we have set $x_f x_b^* = R e^{i\beta t}$, R being a complex constant. For $n > 0$, the stationary state is given by

$$(4Tn + i\gamma\beta) f_n + [Rf_{n-1} - R^*f_{n+1}] = 0. \quad (36)$$

By introducing $Z_n = R^*(f_{n+1}/f_n)$ and solving the equation by means of the continuous fraction method, we obtain the recurrence relation

$$Z_{n-1} = \frac{|R|^2}{Z_n - (4Tn + i\gamma\beta)}. \quad (37)$$

The normalisation condition for ρ gives $2\pi f_0 = 1$, while the atomic contribution for x_b and x_f can be then determined from Eq. (28),

$$\begin{aligned} B_1 &= 2\pi f_{-1} e^{-i\beta t} = \frac{Z_0^*}{R} e^{-i\beta t} \\ B_{-1} &= 2\pi f_1 e^{i\beta t} = \frac{Z_0}{R^*} e^{i\beta t}. \end{aligned} \quad (38)$$

This methods enables us to go beyond the linear analysis determining the dependence of *both* x_f and x_b on the injected field (see Section V), with the only warning that Z_0 has to be computed numerically.

V. PREDICTIONS

In this section we discuss the results derived in Section IV in the large detuning limit for both the Fokker-Planck (see Sec. IV B) and Vlasov (Sec. IV A) models. We first test the validity of the two approaches by comparing them to the experimental results of Ref. [11]. The experiment has been performed with cold ^{85}Rb atoms in a high- Q ring cavity. The atomic parameters are:

$m = 1.4 \cdot 10^{-29} \text{kg}$, $\mathcal{D} = 1.5 \cdot 10^{-29} \text{Cm}^{-1}$ with relaxation rates for atomic polarisation and inversion $2\gamma_{\perp} = \gamma_{\parallel} = 5.9 \text{MHz}$, respectively. Moreover, the atomic frequency is $\omega = 3.77 \times 10^{14} \text{Hz}$ (for the D_1 line), while the input field is shifted 1THz away from it. The beam diameter is $130 \mu\text{m}$. The cavity linewidth is 22kHz , the medium's length $L = 10^{-3} \text{m}$, and the transmissivity $\mathcal{T} = 1.8 \times 10^{-6}$. Since a servo-mechanism continuously adapts the dressed cavity resonance to the frequency of the injected field, we assume $\bar{\Delta} = 0$.

The equilibrium gas temperature, in the absence of collective interaction, is experimentally known with a large uncertainty. We use an estimated value, $\bar{T} \approx 250 \mu\text{K}$, which lies within the range of uncertainty. The second crucial parameter which characterizes the thermalisation is the damping constant, experimentally evaluated in Ref. [11] to be $\gamma = 9$. We will thus use $\gamma = \gamma_c = 9$.




FIG. 3: Bifurcation diagram numerically computed using the Fokker-Planck model. Figs. (a), (b), (c), present respectively the backward field power, P_B , the forward field power, P_F and the detuning between backward and forward field, β , as function of the injected power P_I . All quantities are in physical units. The parameters correspond to those of the experiment [11].

We first report the results of the Fokker-Planck model since it describes more appropriately the molasses dynamics. In Fig. 3a one remarks, from the shape of the backward field power, the existence of a continuous phase transition. The threshold is located at $P_I \approx 0.2 \text{W}$, compatible with the measurements of Ref. [11], which reports backward lasing as occurring for a few Watt of injected power.

The possible underestimate in the threshold value, which we obtain from our approach, can be attributed to two experimental features that cannot be taken into account in a simple way. First, the servo-control which keeps the cavity on resonance by *maximizing the forward transmitted field* [11] cannot be correctly modeled by simply setting $\bar{\Delta} = 0$. Indeed, its operation amounts to a dynamical process which counteracts the birth of the backward emission, since the latter removes energy from the forward field. As a result, one expects the thresh-

old to be displaced towards higher input power values. Second, the spatial modulation imprinted onto the sample by the molasses beam, with a wavelength that differs from that of the collective process (by a factor close to $\sqrt{2}$), introduces a competition between two incompatible length scales. It is likely that this effect should also contribute to raising the critical input power even further.

The detuning value, β , at which the backward field arises (cf. Fig. 3c) also agrees with the observations [11], which report a red-shift around 100kHz. Moreover, the continued fraction expansion discussed in the previous section also allows for a determination of the behavior of the forward field as a function of the input power (cf. Fig. 3b). Below threshold, P_F grows linearly as a function of the injected power, P_I , as no energy is transferred to the backward field. Above threshold, a fraction of the injected field feeds the backward mode, thus leading to a slower growth of P_F .




FIG. 4: Bifurcation diagram numerically computed using the Vlasov model. Figs. (a), (b), (c), present respectively the backward field power, P_B , the forward field power, P_F , and the detuning between backward and forward field, β , as function of the injected power P_I . All quantities are in physical units. The parameters correspond to those of the experiment [11].

Although one does not expect the Vlasov description to closely reproduce the experimental results reported in Ref. [11], it is worth analyzing its predictions; this will help us clarifying the role of the relaxation mechanisms and will sharpen our understanding of the problem. From Fig. 4 one sees that, for the same parameter values used in the Fokker-Planck approach, the threshold is somewhat lower and the red shift in the backward field frequency is smaller as well. Nevertheless, the differences are not so dramatic, since, while the threshold occurs for somewhat lower values of the input power, the frequency shift shows an even somewhat better agreement with the observations [11]. In addition, the uncertainty on some of the experimental parameters is large enough to render a sure discrimination between the modeling based on a Vlasov or on a Fokker-Planck process quite difficult. Be-

cause of the intrinsic paradox, given by the inclusion in the experiment of the optical molasses which ought to exclude the Vlasov description, this result strongly indicates that the current state of the system's characterization, both experimental and theoretical, is not adequate; hence, a large amount of caution in drawing conclusions ought to be exercised.




FIG. 5: Position distribution $\rho(\theta)$, for different values of the injected power P_I , in physical units. In each part of the figure, the dotted line corresponds to the uniform distribution. Parameters correspond to the experimental values of Ref. [11].

In addition to the direct comparison to the experiment, we can also examine some specific predictions coming from each model. The distribution of atomic positions ρ predicted by the Fokker-Planck model is plotted in Fig. 5: the different parts of the figure correspond to increasing values of the injected power P_I (the other parameters remaining unchanged). One can clearly see the rise of the modulation with an increasing role played by higher harmonics.

In the Vlasov description, there is no equivalent regime characterized exclusively by the distribution of θ -values. The probability density $Q(\theta, p)$ for an input intensity




FIG. 6: Joint distribution $Q(\theta, p)$, above threshold. The parameter values are the same as those of the experiment [11] with input power of 480mW.

above threshold is plotted in Fig. 6, where one can indeed appreciate the need to account for both the p and θ dependence in this framework. Nevertheless, one can compute $\rho(\theta)$ by integrating $Q(\theta, p)$ (see Eq. (24)). The results, plotted in Fig. 7, reveal again a qualitative agreement with the Fokker-Planck approach. The larger modulation for the same input power is a consequence of the smaller value of the threshold in the Vlasov description.




FIG. 7: Atomic position distribution $\rho(\theta)$, as function of θ , above threshold. The dotted line corresponds to the uniform distribution, below threshold.

VI. CONCLUSIONS AND PERSPECTIVES

In this paper we have presented a general framework for the description of the *spontaneous* appearance of a *stationary* density grating arising from the interaction between an atomic sample enclosed in a bidirectional ring cavity and an injected, quasi-resonant field. This contribution completes the picture previously sketched out in Refs. [12, 14], where we have shown that the inclusion of a relaxation mechanism for the external degrees of freedom is indispensable for a correct modeling of the interaction, and is responsible for the appearance of a stationary backward field. In the previous work, we focused on the existence of a different kind of phase transition, initiated by a grating in the phase of the atomic polarization. Here, with the help of a detailed numerical analysis we have proven beyond doubt, that a true stationary CARL action – i.e., initiated by the birth of an atomic density modulation – can spontaneously occur. For the investigation, we have chosen to study a bidirectional ring cavity model [13, 15] because of the fuller and more correct description that the latter provides of a real system. In the course of the presentation, we have shown that in certain limits the ring cavity model reduces to the single-pass one [4, 12, 14]; this legitimizes, a posteriori, the use of the latter approach under such conditions.

Following a discussion of the main relaxation mechanisms that one may expect to find in an experiment, we have offered an analytical description of the two most

likely ones on the basis of a probabilistic description, in the large detuning limit. Comparison between our predictions and recent experimental results [11] shows a semi-quantitative agreement, surprisingly, for both the Fokker-Planck and the Vlasov approach. Besides the need for a more accurate experimental characterization of the phenomenon, including a better determination of the parameter values, and systematic measurements of the threshold behavior and of the functional dependence of the backward field power and frequency, additional theoretical work needs to be completed. In particular, a more accurate modeling of the relaxation mechanism represented by the optical molasses needs to be achieved, together with an analysis of the limits of validity of the descriptions proposed for the losses. Finally, it will be useful to investigate the transtion scenario when two mech-

anisms separately described here and in Refs. ([13, 15]) cooperate to induce a backward field; this is likely to occur in the range of smaller detunings when the atomic variables cannot be eliminated.

Acknowledgments

The authors warmly thank Ph.W. Courteille for sharing information on the experiment before publication. MP wishes to acknowledge INOA for financial support, L.M. Narducci and M. Le Bellac for useful discussions. This work has been partial funded by the FIRB-contract n. RBNE01CW3M.001.

-
- [1] C.N. Cohen-Tannoudji, Rev. Mod. Phys. **70**, 707 (1998).
 - [2] J.Guo, P.R.Berman, B.Dubetsky, G.Grynberg, Phys.Rev. A **46**, 1426 (1992).
 - [3] J.Y.Courtois, G.Grynberg, B.Lounis, P.Verkerk, Phys. Rev. Lett. **72**, 3017 (1992).
 - [4] R. Bonifacio et L.De Salvo, Nucl. Instr. Methods Phys. Res. Sec. A **341**, 360 (1994); R. Bonifacio, L. De Salvo, L.M. Narducci, and E.J. D'Angelo, Phys Rev. A **50**, 1716 (1994).
 - [5] P.R. Berman Phys.Rev.A **59**, 585 (1999).
 - [6] L.A. Orozco, H.J. Kimble, et al., Phys. Rev. A **39**, 1235 (1989). See section V. E.
 - [7] G.L. Lippi, G.P. Barozzi, S. Barbay, and J.R. Tredicce, Phys. Rev. Lett. **76**, 2452 (1996).
 - [8] P.R. Hemmer, N.P. Bigelow, D.P. Katz, M.S. Shahriar, L. DeSalvo, and R. Bonifacio, Phys. Rev. Lett. **77**, 1468 (1996).
 - [9] S. Barbay, G. Fabre, and G.L. Lippi, Opt. Commun. **165**, 119 (1999).
 - [10] W.J. Brown, J.R. Gardner, D.J. Gauthier, and R. Vialaseca, Phys. Rev. A **55**, R1601 (1997).
 - [11] D. Kruse, C. von Cube, C. Zimmermann, Ph.W. Courteille, Phys. Rev. Lett., **91**, 183601 (2003).
 - [12] M.Perrin, G.L.Lippi, A.Politi, Phys. Rev. Lett. **86**, 4520 (2001).
 - [13] M. Perrin, Z. Ye, L.M. Narducci, Phys. Rev. A **66**, 043809 (2002).
 - [14] J. Javaloyes, G.L. Lippi, A. Politi, Phys. Rev. A **68**, 033405 (2003).
 - [15] Z. Ye and L.M. Narducci, Phys. Rev. A **63**, 043815 (2001).
 - [16] For instance, the original model [4], derived under the approximation of a constant input field, does not conserve energy even in the absence of dissipation.
 - [17] F. Mitschke, R. Deserno, W. Lange, and J. Mlynek, Phys. Rev. A **33**, 3219 (1986).
 - [18] T. Ackemann and W. Lange, Appl. Phys. B **72**, 21 (2001).
 - [19] R.A. MacDonald and D.H. Tsai, Phys. Rep. **6**, 1 (1978).
 - [20] C. Cohen-Tannoudji, J.Dupont-Roc, G.Grynberg, *Atom-Photon interactions: Basic Processes and Applications*, Wiley (New York, 1992).
 - [21] While the atomic polarization and population can be safely eliminated by following the procedure described in Ref. [14], more questionable is the treatment of position and momentum. However, preliminary investigations indicate that this is possible at least in some parameter ranges.
 - [22] S. Haroche, F. Hartmann, Phys. Rev. A **6**, 1280 (1972).
 - [23] R. Bonifacio and P. Verkerk, Opt. Commun. **124**, 469 (1996).
 - [24] R. Bonifacio and L. DeSalvo, Opt.Comm. **115**, 505 (1995).
 - [25] The net effect of this randomization is to increase the threshold value of the injected field, because of the disorder introduced.
 - [26] An analogous stability analysis was carried out also in Ref. [23], where, however, the possible occurrence of a phase transition was missed because of the additional presence of a probe field. In fact, as is the case of ferromagnetic systems, the presence of an external field destroys the second-order transition.
 - [27] H.Risken, *The Fokker-Planck equation*, Springer Verlag (1984), see section 10.4.
 - [28] S.H. Strogatz, Physica (Amsterdam), **143D**, 1 (2000)